

II. AMENDMENTS TO THE CLAIMS

Please amend the claims as follows:

1. **(Original)** An oral dosage form, comprising
an orally therapeutically effective amount of an opioid agonist, and
an opioid antagonist, the ratio of opioid antagonist to opioid agonist providing a combination
product which is analgesically effective when the combination is administered orally, but which
is aversive in physically dependent human subjects when administered at the same dose or at a
higher dose than the usually prescribed dose of the opioid agonist.
2. **(Original)** The oral dosage form of claim 1, wherein the amount of antagonist
included in the oral dosage form causes an aversive experience in a physically dependent addict
taking about 2-3 times the usually prescribed dose of the opioid.
3. **(Original)** The oral dosage form of claim 1, wherein the opioid agonist is
hydrocodone and the antagonist is naltrexone.
4. **(Original)** The oral dosage form of claim 3, wherein the ratio of naltrexone to
hydrocodone is from about 0.03:1 to about 0.27:1.
5. **(Original)** The oral dosage form of claim 3, wherein the ratio of naltrexone to
hydrocodone is from about 0.05:1 to about 0.20:1.
6. **(Original)** The oral dosage form of claim 1, wherein the opioid agonist is selected
from the group consisting of morphine, hydromorphone, hydrocodone, oxycodone, codeine,
levorphanol, meperidine, methadone, and mixtures thereof.
7. **(Original)** The oral dosage form of claim 1, further comprising an additional non-

opioid drug selected from the group consisting of an NSAID, a COX-2 inhibitor, acetaminophen, aspirin, an NMDA receptor antagonist, a drug that blocks a major intracellular consequence of NMDA-receptor activation, an antitussive, an expectorant, a decongestant, an antihistamine and mixtures thereof.

8. **(Original)** The oral dosage form of claim 1, further comprising one or more pharmaceutically acceptable inert excipients.

9. **(Original)** The oral dosage form of claim 6, wherein said opioid antagonist is selected from the group consisting of naltrexone, naloxone, nalmephene, cyclazocine, levallorphan, and mixtures thereof.

10. **(Original)** The oral dosage form of claim 6, wherein said opioid antagonist is naltrexone.

11. **(Original)** The oral dosage form of claim 1, further comprising a sustained release carrier which imparts sustained release properties to said opioid agonist.

12. **(Original)** The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone and said opioid agonist is oxycodone, wherein the ratio of naltrexone to oxycodone is from about 0.037:1 to about 0.296:1.

13. **(Original)** The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone and said opioid agonist is codeine, wherein the ratio of naltrexone to codeine is from about 0.005:1 to about 0.044:1.

14. **(Original)** The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone and said opioid agonist is hydromorphone, wherein the ratio of naltrexone to

hydromorphone is from about 0.148:1 to about 1.185:1.

15. **(Original)** The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone and said opioid agonist is levorphanol, wherein the ratio of naltrexone to levorphanol is from about 0.278:1 to about 2.222:1.

16. **(Original)** The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone and said opioid agonist is meperidine, wherein the ratio of naltrexone to meperidine is from about 0.0037:1 to about 0.0296:1.

17. **(Original)** The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone and said opioid agonist is methadone, wherein the ratio of naltrexone to methadone is from about 0.056:1 to about 0.444:1.

18. **(Original)** The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone and said opioid agonist is morphine, wherein the ratio of naltrexone to morphine is from about 0.018:1 to about 0.148:1.

19. **(Original)** The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone and said opioid agonist is oxycodone, wherein the ratio of naltrexone to oxycodone is from about 0.056:1 to about 0.222:1.

20. **(Original)** The oral dosage form of claim 1, wherein said opioid antagonist is naltrexone and said opioid agonist is codeine, wherein the ratio of naltrexone to codeine is from about 0.0083:1 to about 0.033:1.

Claims 21 - 35 **(Cancelled)**.